






Analytical Hierarchy Process


- A systematic method for comparing a list of objectives or alternatives
- When used in the systems engineering process, AHP can be a powerful tool for comparing alternative design concepts
- **Reference:** Ernest H. Forman, *Decision by Objectives*,
<http://www.expertchoice.com/dbo/>








- Assume that a set of objectives has been established (VSD, OH), and that we are trying to establish a normalized set of weights to be used when comparing alternatives using these objectives.
- For simplicity, we assume that there are 4 objectives: O_1 , O_2 , O_3 , and O_4 .



- Form a pairwise comparison matrix A , where the number in the i_{th} row and j_{th} column gives the relative importance of O_i as compared with O_j
- Use a 1–9 scale, with
 - $a_{ij} = 1$ if the two objectives are equal in importance
 - $a_{ij} = 3$ if O_i is weakly more important than O_j
 - $a_{ij} = 5$ if O_i is strongly more important than O_j
 - $a_{ij} = 7$ if O_i is very strongly more important than O_j
 - $a_{ij} = 9$ if O_i is absolutely more important than O_j
 - $a_{ij} = 1/3$ if O_j is weakly more important than O_i




- Thus we might arrive at the following matrix:

$$A = \begin{bmatrix} 1 & 1/5 & 1/3 & 1/7 \\ 5 & 1 & 3 & 5 \\ 3 & 1/3 & 1 & 3 \\ 7 & 1/5 & 1/3 & 1 \end{bmatrix} = \begin{bmatrix} 1.000 & 0.200 & 0.333 & 0.143 \\ 5.000 & 1.000 & 3.000 & 5.000 \\ 3.000 & 0.333 & 1.000 & 3.000 \\ 7.000 & 0.200 & 0.333 & 1.000 \end{bmatrix}$$


- To normalize the weights, compute the sum of each column and then divide each column by the corresponding sum
- Using an overbar to denote normalization, we get:

$$\bar{A} = \begin{bmatrix} 0.063 & 0.115 & 0.071 & 0.016 \\ 0.313 & 0.577 & 0.643 & 0.547 \\ 0.188 & 0.192 & 0.214 & 0.328 \\ 0.438 & 0.115 & 0.071 & 0.109 \end{bmatrix}$$



$$\bar{A} = \begin{bmatrix} 0.063 & 0.115 & 0.071 & 0.016 \\ 0.313 & 0.577 & 0.643 & 0.547 \\ 0.188 & 0.192 & 0.214 & 0.328 \\ 0.438 & 0.115 & 0.071 & 0.109 \end{bmatrix}$$

- The numbers in the second row are generally larger than the rest of the numbers, except for the case of column 1
- Indicates some inconsistency in the comparisons used in the original matrix
- Ideally, the 4 normalized columns would all be identical if the pairwise comparisons were consistent
- In practice, one can compute a consistency measure using the eigenvalues of the normalized comparison matrix.



- The next step is to compute the average values of each row and use these as the weights in the Objective Hierarchy
- For this example, the weights would be:

$$w = [0.066 \quad 0.520 \quad 0.231 \quad 0.183]$$

- Note that by construction, $\sum_{i=1}^4 w_i = 1$.
- These weights would be used in summing the measures as required in the evaluation of the Objective Hierarchy.